

What You Need To Know About H1N1 (swine flu) & Keeping Your Family Healthy

By: Kara Gabriele, Ph.D.

Information taken from the Center for Disease Control Website: www.cdc.gov

What is novel H1N1 (swine flu)?

Novel H1N1 (referred to as “swine flu” early on) is a new influenza virus causing illness in people. This new virus was first detected in people in the United States in April 2009. This virus is spreading from person-to-person worldwide, probably in much the same way that regular seasonal influenza viruses spread. On June 11, 2009, the World Health Organization (WHO) signaled that a pandemic of novel H1N1 flu was underway.

How does novel H1N1 virus spread?

Flu viruses are spread mainly from person to person through coughing or sneezing by people with influenza. Sometimes people may become infected by touching something – such as a surface or object – with flu viruses on it and then touching their mouth or nose.

What are the signs and symptoms of this virus in people?

The symptoms of novel H1N1 flu virus in people include fever, cough, sore throat, runny or stuffy nose, body aches, headache, chills and fatigue. A significant number of people who have been infected with this virus also have reported diarrhea and vomiting.

How severe is illness associated with novel H1N1 flu virus?

Illness with the new H1N1 virus has ranged from mild to severe. While most people who have been sick have recovered without needing medical treatment, hospitalizations and deaths from infection with this virus have occurred. In seasonal flu, certain people are at “high risk” of serious complications. This includes people 65 years and older, children younger than five years old, pregnant women, and people of any age with certain chronic medical conditions.

How does novel H1N1 flu compare to seasonal flu in terms of its severity and infection rates?

Seasonal influenza can cause mild to severe illness, and at times can lead to death. Each year, in the United States, on average 36,000 people die from flu-related complications and more than 200,000 people are hospitalized from flu-related causes. Of those hospitalized, 20,000 are children younger than 5 years old. Over 90% of deaths and about 60 percent of hospitalization occur in people older than 65.

How long can an infected person spread this virus to others?

People infected with seasonal and novel H1N1 flu viruses may be able to infect others from 1 day before getting sick to 5 to 7 days after. This can be longer in some people, especially children and people with weakened immune systems and in people infected with the new H1N1 virus.

Prevention & Treatment

What can I do to protect myself from getting sick?

There is no vaccine available right now to protect against novel H1N1 virus. However, a novel H1N1 vaccine is currently in production and may be ready for the public in the fall. As always, a vaccine will be available to protect against [seasonal influenza](#). There are everyday actions that can help prevent the spread of germs that cause respiratory illnesses like influenza.

Take these everyday steps to protect your health:

- Cover your nose and mouth with a tissue when you cough or sneeze. Throw the tissue in the trash after you use it.
- Wash your hands often with soap and water for a minimum of 20 seconds, especially after you cough or sneeze. [Alcohol-based hand cleaners*](#) are also effective.
- Avoid touching your eyes, nose or mouth. Germs spread this way.
- Try to avoid close contact with sick people.
- If you are sick with flu-like illness, [CDC recommends that you stay home for at least 24 hours after your fever is gone](#) except to get medical care or for other necessities. (Your fever should be gone without the use of a fever-reducing medicine.) Keep away from others as much as possible to keep from making others sick.

Other important actions that you can take are:

- Follow public health advice regarding school closures, avoiding crowds and other social distancing measures.
- Be prepared in case you get sick and need to stay home for a week or so; a supply of over-the-counter medicines, [alcohol-based hand rubs,*](#) tissues and other related items might could be useful and help avoid the need to make trips out in public while you are sick and contagious

What is the best way to keep from spreading the virus through coughing or sneezing?

If you are sick with flu-like illness, [CDC recommends that you stay home for at least 24 hours after your fever is gone](#) except to get medical care or for other necessities. (Your fever should be gone without the use of a fever-reducing medicine.) Keep away from others as much as possible. Cover your mouth and nose with a tissue when coughing or sneezing. Put your used tissue in the waste basket. Then, clean your hands, and do so every time you cough or sneeze.

What is the best technique for washing my hands to avoid getting the flu?

Washing your hands often will help protect you from germs. Wash with soap and water or clean with [alcohol-based hand cleaner*](#). CDC recommends that when you wash your hands -- with soap and warm water -- that you wash for **15 to 20 seconds**.

What should I do if I get sick?

If you live in areas where people have been identified with novel H1N1 flu and become ill with influenza-like symptoms, including fever, body aches, runny or stuffy nose, sore throat, nausea, or vomiting or diarrhea, you should stay home and avoid contact with other people. [CDC recommends that you stay home for at least 24 hours after your fever is gone](#) except to get medical care or for other necessities. (Your fever should be gone without the use of a fever-reducing medicine.) Stay away from others as much as possible to keep from making others sick. Staying at home means that you should not leave your home except to seek medical care. This means avoiding normal activities, including work, school, travel, shopping, social events, and public gatherings.

If you become ill and experience any of the following warning signs, seek emergency medical care.

In children, emergency warning signs that need urgent medical attention include:

- Fast breathing or trouble breathing
- Bluish or gray skin color
- Not drinking enough fluids
- Severe or persistent vomiting
- Not waking up or not interacting
- Being so irritable that the child does not want to be held
- Flu-like symptoms improve but then return with fever and worse cough

In adults, emergency warning signs that need urgent medical attention include:

- Difficulty breathing or shortness of breath
- Pain or pressure in the chest or abdomen
- Sudden dizziness
- Confusion
- Severe or persistent vomiting
- Flu-like symptoms improve but then return with fever and worse cough

Are there medicines to treat novel H1N1 infection?

Yes. CDC recommends the use of oseltamivir or zanamivir for the treatment and/or prevention of infection with novel H1N1 flu virus. Antiviral drugs are prescription medicines (pills, liquid or an inhaled powder) that fight against the flu by keeping flu viruses from reproducing in your body. If you get sick, antiviral drugs can make your illness milder and make you feel better faster. They may also prevent serious flu complications.

Contamination & Cleaning

How long can influenza virus remain viable on objects (such as books and doorknobs)

Studies have shown that influenza virus can survive on environmental surfaces and can infect a person for 2 to 8 hours after being deposited on the surface.

What kills influenza virus?

Influenza virus is destroyed by heat (167-212°F [75-100°C]). In addition, several chemical germicides, including chlorine, hydrogen peroxide, detergents (soap), iodophors (iodine-based antiseptics), and alcohols are effective against human influenza viruses if used in proper concentration for a sufficient length of time. For example, wipes or gels with alcohol in them can be used to clean hands. The gels should be rubbed into hands until they are dry.

***What if soap and water are not available and alcohol-based products are not allowed in my facility?**

Though the scientific evidence is not as extensive as that on hand washing and alcohol-based sanitizers, other hand sanitizers that do not contain alcohol may be useful for killing flu germs on hands.

What surfaces are most likely to be sources of contamination?

Germs can be spread when a person touches something that is contaminated with germs and then touches his or her eyes, nose, or mouth. Droplets from a cough or sneeze of an infected person move through the air. Germs can be spread when a person touches respiratory droplets from another person on a surface like a desk, for example, and then touches their own eyes, mouth or nose before washing their hands.